AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

- 1. 5. (Cancelled)
- 6. (Currently Amended) A method of treatment comprising administering to a patient suffering from, or at a risk of developing, diagnosed with fibromyalgia about 15 to about 360 mg AAT per kg patient body mass, and repeating the administration at least once with a periodicity of between 3 and 31 days.
- 7. (Original) Method, according to claim 6, wherein AAT is administered at a dose of between 25 and 6 mg/kg every week or multiples of these quantities adjusted according to the time interval foreseen until the next dose, in a proportional manner.
- 8. (Currently Amended) A method of treatment comprising administering sufficient AAT to a patient suffering from, or at a risk of developing, diagnosed with fibromyalgia as will increase the patient's AAT level, even by about 100% over basal level at about 7 days following administration and repeating the administration at least once at about 3 to 31 days.
- 9. (Original) Method, according to claim 8, wherein the administration of AAT is repeated at least once at about 7 to 21 days.
- 10. (Currently Amended) A method of treatment comprising administering to a patient suffering from, or at a risk of developing, diagnosed with

fibromyalgia a quantity of AAT as will increase serum AAT levels, even to about 5-fold greater than basal levels at 24 hours following administration.

- 11. (Original) Method, according to claim 10, wherein the administration is repeated one or more times at about 3 to 31 days.
- 12. (Original) Method, according to claim 10, wherein the administration is repeated one or more times at about 7 to 21 days.
- 13. (Currently Amended) Method, according to claim 10, wherein a quantity of AAT is administered as will increase serum AAT levels even about 8-fold over basal levels at 24 hours following administration, and the administration is repeated one or more times at about 7 to 21 days.
- 14. (Currently Amended) A method of treating a patient suffering from, or at a risk of developing, diagnosed with fibromyalgia comprising administering to said patient by parenteral infusion about 15 to 360 mg AAT per kg patient body mass, and repeating said infusion at least once at 3 to 31 days to effect a diminishment of the symptoms of fibromyalgia in said patient.
- 15. (Original) Method, according to claim 14, wherein the dose of AAT is between 25 and 60 mg AAT/kg body mass, and the infusion is repeated at least once at about 7 to 21 days.
- 16. (Original) Method, according to claim 14 or 15, wherein the administration is carried out by intravenous infusion.